



**Gramin Krishi Mausam Sewa**  
**District Level Agromet Advisory Bulletin**  
**ICAR Complex Branch**  
 Umiam, Meghalaya



## Agromet Advisory Bulletin

Date : 29-07-2022

Weather Forecast of District THOUBAL(Manipur) Issued On : 2022-07-29(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-30	2022-07-31	2022-08-01	2022-08-02	2022-08-03
Rainfall(mm)	3.0	6.0	7.0	6.0	14.0
Tmax(°C)	33.0	32.0	31.0	30.0	30.0
Tmin(°C)	23.0	23.0	22.0	22.0	22.0
RH-I(%)	98	97	98	98	96
RH-II(%)	53	65	61	71	67
Wind Speed(kmph)	7.2	7.2	7.0	6.5	5.1
Wind Direction(Degree)	288	291	291	288	290
Cloud Cover(Octa)	8	8	8	8	8

### Weather Summary/Alert:

There is likely to have light to moderate rains during Saturday to Wednesday with max. temp. 30-33 oC & mini. temp. 22-23 oC. Relative humidity may reach upto 98 % (max) & 71 % (min). The average wind speed may be 5-7 km/hr. The extended range forecast indicates that during 3 to 9 August, 2022 rainfall is likely to be normal, minimum temperature and maximum temperature. are likely to be moderately above normal over the state.

### General Advisory:

1. Masks and Frequent Hygiene Measures: Wearing a mask during work time and afterwards helps reduce the spread of COVID-19. Everyone should wear a mask correctly. It should cover your nose and mouth and be changed frequently. Everyone should frequently wash their hands or use a hand sanitizer with at least 60% alcohol. 2. Hygiene and Precaution: When coming back from the fields, wear a mask, wash hands, use hand sanitizer, and wash worker's clothes separately. Cover your mouth and nose with a tissue when you cough or sneeze. 3. Clean and Safe Household: People should stay at least 6 feet apart in shared areas. Don't share your tools and utensils. Keep all frequently touched surfaces clean. Maintain proper ventilation by opening windows and doors. 4. People at High Risk: If you or anyone else in your household is at high risk (older adults and people with illnesses), you should take additional physical distancing and hygiene precautions. 5. If Anyone Is Sick: If you or anyone else in household has symptoms of COVID-19 like fever/ cough, they should isolate and not use shared rooms/ utensils/ areas, if available, use separate toilet. Everyone in the household should wear masks. 6. Medical Assistance: Ask your health care provider about treatment for COVID-19 and how to get a vaccine. COVID-19 vaccines are safe and available, regardless of your immigration or health insurance status. 7. Get Vaccinated: COVID-19 vaccines are safe. Everyone should get vaccinated to protect ourselves, our families, and friends. Get 1st dose of Covid-19 vaccine and after 12-16 weeks get 2nd dose of it. 8. Clean the work tools properly after the work. 9. Consider cancelling all public or nonessential group activities and events. 10. Taking care of ourselves only can help in reducing the spread of Covid-19.

### SMS Advisory:

In those fields where rice was not cultivated due to lack of water, if water is available, RC Maniphou 7, RC Maniphou 12 and RC Maniphou 13 varieties may be sown till August.

**Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
RICE	Kharif rice: (nursery raising for late transplanting to main land preparation and transplanting) - In those fields where rice was not cultivated due to lack of water, if water is available, RC Maniphou 7, RC Maniphou 12 and RC Maniphou 13 varieties may be sown till August. - Water level just to submerge the soil surface in the kharif-rice field may be maintained. - For Direct seeding with drum seeder, soak well filled disease free seeds of rice and incubate to favour germination. Use slightly germinated seeds. Treat seed treatment with Humicil @ 5ml/kg seed. Puddle the field properly and level it. Drain out water 24 hrs before sowing. - Use Blue Sticky Cards@ 4 cards/ sangam or spray Chloropyriphos @ 1ml/ litre water against Thrips. - Apply Butachlor @25kg/ha within one week of transplanting rice for weed control. Maintain water level to submerge the soil surface. Regularly monitor the water level in the crop field.
GINGER	Ginger/ Turmeric (rhizome formation): - Prepare to avoid excess rain water logging in the crop field. - To prevent root/ rhizome rot drench the soil with Mancozeb @3g/ L water or Metalaxyl @ 1.25 g/ L water. - To prevent bacterial wilt, drench the soil with Copper oxychloride @3g/ L water immediately after disease is seen in the field. - Weeding and earthing up may be done to enhance root development after rains. Repeat green leaves mulching after inter-culture operations @ 7.5t/ ha. - To prevent Leaf Spot disease, apply Bordeaux mixture (5:5:50) after 4 – 5 weeks of sowing or Captan @ 1.5 g/l of water after rains. - Short duration French Bean may sown as inter-crop.
SOYABEAN	Soybean/ groundnut: (vegetative to flowering) - Avoid high weed population by weeding when the soil condition permits after rains. Construct proper drainage system in and around the crop field to avoid water stagnation especially during rainy days. - Weed control must be strictly done in the field upto 60 days after sowing for healthy growth of the crops. - If incidence of hairy caterpillar is observed then weeding may be done and spray Dichlorovos 76% EC @ 1ml/ lit of water along the field bunds by making trenches.
MAIZE	Maize (vegetative to flowering) - Avoid weed in the crop field. - Regularly monitor insect pest infestation in the crop field.

**Horticulture Specific Advisory:**

Horticulture(Varieties)	Horticulture Specific Advisory
COLOCASIA/ TARO	Colocasia: (vegetative to corm formation) - If the weed population is high then weeding may be done. - As prophylactic measure against leaf blight, spray Redomil @ 1.5g/ litre of water or Mancozeb @ 2gm/lit of water twice or thrice at 10 days interval after the wet spell.
CABBAGE	Cole Crops (Cauliflower/ Cabbage/ Broccoli) (nursery raising to transplanting): - Raise nursery of cole crops under low cost polyhouse. Seed rate is 600-750g/ha for early variety. - Ready seedlings of cole crops may be transplanted.
ORANGE	For grown-up trees, apply 450 g Urea, 1000 g SSP, 400 g MOP and 50 g Borax per plant. Spray Ridomil @ 2.5 g/lit or Carbandazim @ 10 g/15 litres of water against disease and Imidachloprid @ 5 ml/10 lit water against insect pest. - If lichen (fungus + algae) is observed, then scrap out the affected area and paint with Bordeaux paste or Benlate. Prune the disease infected twigs and burn them outside the farm area. Spray Copper oxychloride @ 3ml/lit to control fungal disease.
LEMON	For grown-up trees, apply 400g Urea, 550g SSP, 250g MOP and 50g Borax per plant. Spray Ridomil @ 2.5 g/lit or Carbandazim @ 10 g/15 litres of water against disease and Imidachloprid @ 5 ml/10 lit water against insect pest. - If lichen (fungus + algae) is observed, then scrap out the affected area and paint with Bordeaux paste or Benlate. Prune the disease infected twigs and burn them outside the farm area. Spray Copper oxychloride @ 3ml/lit to control fungal disease.

**Live Stock Specific Advisory:**

Live Stock(Varieties)	Live Stock Specific Advisory
-----------------------	------------------------------

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Cattle: - Prepare to avoid rain water logging in and around the shed. - Protect animals from rains and provide clean drinking water. - Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal. - Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated. - De-worm with Albomac suspension@60-90ml for 300kg body weight or 20-30ml for 100 kg body weight. - Provide clean and safe drinking water during rainy days.
PIG	Pig: - Prepare to avoid rain water logging in and around the shed. Provide clean drinking water - Vaccinate pigs against African Swine Fever. - Band import of pigs from neighboring states/ country. - Rinsing with sodium hypochlorite (200 ppm chlorine) and Ethyl alcohol (70 % in water) of skin may be done. - Check on availability and quality of feeds and water; give steamed/boiled feeds, provision may be made for balanced ration with high fiber with leguminous feed and complete stoppage on the importation on the hay, straw and vegetables from the outbreak areas. - Slaughts of animals/ porks processing must be totally banned in and around the outbreak places and neighbouring villages too.

#### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
CARP	Fish (Nursery Pond for Common Carp and Grass Carp) - Treat one hectare pond with a mixture of 100kg Common Salt + 10 kg Garlic paste soaked overnight. - Dykes/ bunds around the fish ponds may be raised high and encircle with happa net to avoid escape of fishes from pond. - Harvest rain water and treat before using it for fish culture. - Prepare fish nursery ponds of 1.5m depth. Maintain water depth of 1m in the pond. - Apply lime @300kg/ha. pH of pond water should be maintained at 7.0-7.5 and apply manure with a slurry of RCD @5000kg/ha. - Stocking of 5-6 million spawns per hectare pond may be done. Later thinning may be done when required. - Well balanced feeds should be provided 2-3 times a day @ 5-10% body weight/day

#### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Give Amprolium @ 2 g/ litre of drinking water for 5-7 days against Coccidiosis. Provide clean drinking water during rainy days. Check for leakage in the sheds. Proper care must be taken to avoid flies and mosquitoes. Monitor regularly the health status of the birds. Treatment must be provided after consulting experts at earliest possible. Add sanitizer in drinking water during rainy days.